

FUN FACTS TO KNOW AND TELL

$$\begin{aligned} I_n &\equiv \int_0^\infty dx \frac{x^{n-1}}{e^x - 1} = \Gamma(n)\zeta(n), & \int_0^\infty dx \frac{x^{n-1}}{e^x + 1} &= \Gamma(n)\zeta(n) \left[1 - (1/2)^{n-1}\right], \\ \zeta(n) &\equiv \sum_{m=1}^\infty m^{-n}, & \Gamma(n) &\equiv (n-1)!, \\ \zeta(3/2) &= 2.612375\dots, & \zeta(2) &= \frac{\pi^2}{6}, & \zeta(3) &= 1.20205\dots, & \zeta(4) &= \frac{\pi^4}{90}, \\ & \int_{-\infty}^\infty dx e^{-x^2/2} = \sqrt{2\pi}, & \int_0^\infty dx x^n e^{-x} &= n! \end{aligned}$$

LONG ANSWER SECTION

1. (10 pts) Beginning with the fundamental thermodynamic relation,

$$TdS = dE + PdV - \mu dN,$$

derive the Maxwell relation

$$\left. \frac{\partial S}{\partial P} \right|_{T,\mu} = - \left. \frac{\partial V}{\partial T} \right|_{\mu,P}.$$

2. (10 pts) Consider a non-relativistic particle in a one-dimensional potential

$$V(x) = \frac{A}{6}x^6.$$

Using some combination of the equipartition, generalized equipartition and/or virial theorems, find the thermal average $\langle x^6 \rangle$ in terms of T and A .

3. A one-dimensional gas of non-relativistic spin-1/2 fermions of mass m confined within a length L is thermalized according to a chemical potential μ and temperature T , i.e., the phase space occupancy is

$$f(p) = \frac{e^{-(E_p - \mu)/T}}{1 + e^{-(E_p - \mu)/T}}.$$

Originally, the temperature is $T = 0$ and the chemical potential is μ .

- (a) (5 pts) In terms of the m , L and μ find the average number of particles N when $T = 0$.
- (b) (5 pts) In terms of m and L , find the single-particle density of states D (number of states per energy) as a function of the single-particle energy ϵ .
- (c) (10 pts) Assuming μ is held constant while the temperature is slightly raised, find the change in the average number of particles to second order in the temperature. Express your answer in the form,

$$N = N_0 + AT + BT^2,$$

solving for A and B in terms of D and $dD/d\epsilon$ evaluated at the Fermi surface.

SECRET STUDENT NUMBER: 60

EXTRA SPACE FOR WORKING #3.

SHORT ANSWER SECTION

4. (2 pts each) Consider three single particle levels, $-\epsilon$, 0 , and ϵ , which are populated by two indistinguishable spin-zero bosons. Let the system be thermalized at temperature T .
- (a) What is the average total energy when $T = 0$? _____
- (b) What is the total entropy when $T = 0$? _____
- (c) What is the average total energy when $T = \infty$? _____
- (d) What is the total entropy when $T = \infty$? _____
5. (3 pts each) Consider a two-dimensional square lattice of coupled three-dimensional oscillators that supports both longitudinal and transverse modes with the same speed of sound c_s . Assume the lattice is one atom thick in the z direction and infinitely long in the x and y direction. Let C/N refer to the specific heat per oscillator.
- (a) As $T \rightarrow 0$, the specific heat from phonons behaves as $C \sim T^n$. What is n ? _____
- (b) What is C/N as $T \rightarrow \infty$? _____
6. (2 pts each) Consider a one-dimensional Ising model at temperature $T > 0$. Label each of the following as true or false.
- (a) In the exact solution there is no phase transition. _____
- (b) In the mean-field solution there is no phase transition. _____
- (c) In the mean-field solution, the critical exponents are the same as they would be for a two-dimensional model. _____
7. (1 pt each) For the following choose between *maximize or minimize* and between S =entropy, E =energy, F =Helmoltz free energy, P =pressure, or G =Gibbs free energy. Circle your choices.
- (a) For a thermalized system at fixed energy, volume and particle number, the system would adjust any order parameter to minimize / maximize the thermodynamic quantity S, E, F, P, G .
- (b) For a thermalized system at fixed volume, temperature, and particle number, the system would adjust any order parameter to minimize / maximize the thermodynamic quantity S, E, F, P, G .
- (c) For a thermalized system at fixed temperature, particle number and pressure, the system would adjust any order parameter to minimize / maximize the thermodynamic quantity S, E, F, P, G .
- (d) For a thermalized system at fixed volume, temperature, and chemical potential, the system would adjust any order parameter to minimize / maximize the thermodynamic quantity S, E, F, P, G .
8. (2 pts each) Pick the appropriate number of dimensions d for each of the following:
- (a) For Bose condensation of a non-relativistic gas to occur, d must be greater than _____.
- (b) According to the Ginzburg criteria, mean field theories will be valid near T_c for d greater than _____.

9. (1 pt each) Assume the free energy density has the form,

$$f(x, T) = \frac{A}{2}m(x)^2 + \frac{B}{4}m(x)^4 + \frac{\kappa}{2}(\nabla m)^2,$$

where $m(x)$ is the magnetization density, and A , B and κ are functions of the temperature T . For each of the following quantities, choose among *ZERO*, *INFINITE* and *FINITE* for what values these quantities approach as $T \rightarrow T_c$ in a standard Ginzburg-Landau picture of a phase transition. Circle your answers.

- A [*ZERO*, *INFINITE*, *FINITE*].
 - B [*ZERO*, *INFINITE*, *FINITE*].
 - κ [*ZERO*, *INFINITE*, *FINITE*].
 - ξ (the correlation length) [*ZERO*, *INFINITE*, *FINITE*].
 - $\Delta F/A$ (the surface free energy for the interface between positive and negative m domains below T_c) [*ZERO*, *INFINITE*, *FINITE*].
 - $\langle F/V \rangle$ (thermal averaged free energy density) [*ZERO*, *INFINITE*, *FINITE*].
 - C_V (the specific heat) [*ZERO*, *INFINITE*, *FINITE*].
10. (1 pt each) Graph several isotherms on a P vs. V graph illustrating the characteristics of a liquid gas phase transition. The graph should include:
- An isotherm with $T > T_c$.
 - An isotherm with $T = T_c$.
 - An isotherm with $T < T_c$.
 - Label the critical point.
 - For the isotherm with $T < T_c$, label the coexistence points.

